

Sum rules for leading and subleading form factors in Heavy Quark Effective Theory using the non-forward amplitude

F. Jugeau

Instituto de Física Corpuscular, Valencia, Spain

A. Le Yaouanc, L. Oliver¹ and J.-C. Raynal

Laboratoire de Physique Théorique (UMR 8627 CNRS)

Université de Paris Sud-XI, Bâtiment 210, 91405 Orsay cedex, France

Abstract

Within the OPE, we formulate new sum rules in Heavy Quark Effective Theory in the heavy quark limit and at order $1/m_Q$, using the non-forward amplitude. In the heavy quark limit, these sum rules imply that the elastic Isgur-Wise function $\xi(w)$ is an alternate series in powers of $(w - 1)$. Moreover, one gets that the n -th derivative of $\xi(w)$ at $w = 1$ can be bounded by the $(n - 1)$ -th one, and the absolute lower bound for the n -th derivative $(-1)^n \xi^{(n)}(1) \geq \frac{(2n+1)!!}{2^{2n}}$. Moreover, for the curvature we find $\xi''(1) \geq \frac{1}{5}[4\rho^2 + 3(\rho^2)^2]$ where $\rho^2 = -\xi'(1)$. These results are consistent with the dispersive bounds, and they strongly reduce the allowed region of the latter for $\xi(w)$. The method is extended to the subleading quantities in $1/m_Q$. Concerning the perturbations of the Current, we derive new simple relations between the functions $\xi_3(w)$ and $\bar{\Lambda}\xi(w)$ and the sums $\sum_n \Delta E_j^{(n)} \tau_j^{(n)}(1) \tau_j^{(n)}(w)$ ($j = \frac{1}{2}, \frac{3}{2}$), that involve leading quantities, Isgur-Wise functions $\tau_j^{(n)}(w)$ and level spacings $\Delta E_j^{(n)}$. Our results follow because the non-forward amplitude depends on three variables $(w_i, w_f, w_{if}) = (v_i \cdot v', v_f \cdot v', v_i \cdot v_f)$, and we consider the zero recoil frontier $(w, 1, w)$ where only a finite number of j^P states contribute $(\frac{1}{2}^+, \frac{3}{2}^+)$. We also obtain new sum rules involving the elastic subleading form factors $\chi_i(w)$ ($i = 1, 2, 3$) at order $1/m_Q$ that originate from the $\mathcal{L}_{kin}$ and $\mathcal{L}_{mag}$ perturbations of the Lagrangian. To the sum rules contribute only the same intermediate states $(j^P, J^P) = (\frac{1}{2}^-, 1^-), (\frac{3}{2}^-, 1^-)$ that enter in the $1/m_Q^2$ corrections of the axial form factor $h_{A_1}(w)$ at zero recoil. This allows to obtain a lower bound on $-\delta_{1/m^2}^{(A_1)}$ in terms of the $\chi_i(w)$ and the shape of the elastic IW function $\xi(w)$. An important theoretical implication is that $\chi'_1(1)$, $\chi'_2(1)$ and $\chi'_3(1)$ ($\chi_1(1) = \chi_3(1) = 0$ from Luke theorem) must vanish when the slope and the curvature attain their lowest values $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$. These constraints should be taken into account in the exclusive determination of $|V_{cb}|$.

LPT Orsay 05-74 (October 2005)

International Workshop on Quantum Chromodynamics : Theory and Experiment, Conversano (Bari, Italy), 16-20 June 2005

¹Speaker at the conference

We will expose the main results that we have obtained in Heavy Quark Effect Theory using the non-forward amplitude and the Operator Product Expansion. We will first examine results in the heavy quark limit on the shape of the Isgur-Wise function. The method is then generalized to the study of the $1/m_Q$ perturbations, that are of two types, namely perturbations of the Current, and perturbations of the Lagrangian. This is a somewhat longer version of the talk than the one that will appear in the Proceedings.

1 Heavy quark limit

In the leading order of the heavy quark expansion of QCD, Bjorken sum rule (SR) [1] relates the slope of the elastic Isgur-Wise (IW) function $\xi(w)$, to the IW functions of the transitions between the ground state and the $j^P = \frac{1}{2}^+, \frac{3}{2}^+$ excited states, $\tau_{1/2}^{(n)}(w)$, $\tau_{3/2}^{(n)}(w)$, at zero recoil $w = 1$ (n is a radial quantum number). This SR leads to the lower bound $-\xi'(1) = \rho^2 \geq \frac{1}{4}$. Recently, a new SR was formulated by Uraltsev in the heavy quark limit [2] involving also $\tau_{1/2}^{(n)}(1)$, $\tau_{3/2}^{(n)}(1)$, that implies, combined with Bjorken SR, the much stronger lower bound $\rho^2 \geq \frac{3}{4}$, a result that came as a big surprise. In ref. [3], in order to make a systematic study in the heavy quark limit of QCD, we have developed a manifestly covariant formalism within the Operator Product Expansion (OPE). We did recover Uraltsev SR plus a new class of SR. Making a natural physical assumption, this new class of SR implies the bound $\sigma^2 \geq \frac{5}{4}\rho^2$ where σ^2 is the curvature of the IW function. Using this formalism including the whole tower of excited states j^P , we have recovered rigorously the bound $\sigma^2 \geq \frac{5}{4}\rho^2$ plus generalizations that extend it to all the derivatives of the IW function $\xi(w)$ at zero recoil, that is shown to be an alternate series in powers of $(w - 1)$.

Using the OPE and the trace formalism in the heavy quark limit, different initial and final four-velocities v_i and v_f , and heavy quark currents, where Γ_1 and Γ_2 are arbitrary Dirac matrices $J_1 = \bar{h}_{v'}^{(c)} \Gamma_1 h_{v_i}^{(b)}$, $J_2 = \bar{h}_{v_f}^{(b)} \Gamma_2 h_{v'}^{(c)}$, the following sum rule can be written [5] :

$$\left\{ \sum_{D=P,V} \sum_n Tr \left[\bar{\mathcal{B}}_f(v_f) \bar{\Gamma}_2 \mathcal{D}^{(n)}(v') \right] Tr \left[\bar{\mathcal{D}}^{(n)}(v') \Gamma_1 \mathcal{B}_i(v_i) \right] \xi^{(n)}(w_i) \xi^{(n)}(w_f) \right. \\ \left. + \text{Other excited states} \right\} = -2\xi(w_{if}) Tr \left[\bar{\mathcal{B}}_f(v_f) \bar{\Gamma}_2 P'_+ \Gamma_1 \mathcal{B}_i(v_i) \right] . \quad (1)$$

In this formula v' is the intermediate meson four-velocity, $P'_+ = \frac{1}{2}(1 + \not{v}')$ comes from the residue of the positive energy part of the c -quark propagator, $\xi(w_{if})$ is the elastic Isgur-Wise function that appears because one assumes $v_i \neq v_f$. $\mathcal{B}_i$ and $\mathcal{B}_f$ are the 4×4 matrices of the ground state B or B^* mesons and $\mathcal{D}^{(n)}$ those of all possible ground state or excited state D mesons coupled to B_i and B_f through the currents. In (1) we have made explicit the $j = \frac{1}{2}^-$ D and D^* mesons and their radial excitations of quantum number n . The explicit contribution of the other excited states is written below.

The variables w_i , w_f and w_{if} are defined as $w_i = v_i \cdot v'$, $w_f = v_f \cdot v'$, $w_{if} = v_i \cdot v_f$. The domain of (w_i, w_f, w_{if}) is [3]

$$\begin{aligned} w_i, w_f &\geq 1 \\ w_i w_f - \sqrt{(w_i^2 - 1)(w_f^2 - 1)} &\leq w_{if} \\ &\leq w_i w_f + \sqrt{(w_i^2 - 1)(w_f^2 - 1)}. \end{aligned} \quad (2)$$

The SR (1) writes $L(w_i, w_f, w_{if}) = R(w_i, w_f, w_{if})$, where $L(w_i, w_f, w_{if})$ is the sum over the intermediate charmed states and $R(w_i, w_f, w_{if})$ is the OPE side. Within the domain (2) one can derive relatively to any of the variables w_i , w_f and w_{if} and obtain different SR taking different limits to the frontiers of the domain.

As in ref. [3] [4], we choose as initial and final states the B meson $\mathcal{B}_i(v_i) = P_{i+}(-\gamma_5)$, $\mathcal{B}_f(v_f) = P_{f+}(-\gamma_5)$ and vector or axial currents projected along the v_i and v_f four-velocities

$$J_1 = \bar{h}_{v'}^{(c)} \not{v}_i h_{v_i}^{(b)}, \quad J_2 = \bar{h}_{v_f}^{(b)} \not{v}_f h_{v'}^{(c)} \quad (3)$$

we obtain SR (1) with the sum of all excited states j^P in a compact form :

$$\begin{aligned} (w_i + 1)(w_f + 1) \sum_{\ell \geq 0} \frac{\ell + 1}{2\ell + 1} S_\ell(w_i, w_f, w_{if}) \sum_n \tau_{\ell+1/2}^{(\ell)(n)}(w_i) \tau_{\ell+1/2}^{(\ell)(n)}(w_f) \\ + \sum_{\ell \geq 1} S_\ell(w_i, w_f, w_{if}) \sum_n \tau_{\ell-1/2}^{(\ell)(n)}(w_i) \tau_{\ell-1/2}^{(\ell)(n)}(w_f) = (1 + w_i + w_f + w_{if}) \xi(w_{if}) \end{aligned} \quad (4)$$

We get, choosing instead the axial currents,

$$J_1 = \bar{h}_{v'}^{(c)} \not{v}_i \gamma_5 h_{v_i}^{(b)}, \quad J_2 = \bar{h}_{v_f}^{(b)} \not{v}_f \gamma_5 h_{v'}^{(c)}, \quad (5)$$

$$\sum_{\ell \geq 0} S_{\ell+1}(w_i, w_f, w_{if}) \sum_n \tau_{\ell+1/2}^{(\ell)(n)}(w_i) \tau_{\ell+1/2}^{(\ell)(n)}(w_f) + (w_i - 1)(w_f - 1)$$

$$\sum_{\ell \geq 1} \frac{\ell}{2\ell - 1} S_{\ell-1}(w_i, w_f, w_{if}) \sum_n \tau_{\ell-1/2}^{(\ell)(n)}(w_i) \tau_{\ell-1/2}^{(\ell)(n)}(w_f) = -(1-w_i-w_f+w_{if})\xi(w_{if}) \quad (6)$$

Following the formulation of heavy-light states for arbitrary j^P given by Falk [5], we have defined in ref. [3] the IW functions $\tau_{\ell+1/2}^{(\ell)(n)}(w)$ and $\tau_{\ell-1/2}^{(\ell)(n)}(w)$, ℓ and $j = \ell \pm \frac{1}{2}$ being the orbital and total angular momentum of the light cloud.

In (3) and (5) S_n is given by

$$S_n = v_{i\nu_1} \cdots v_{i\nu_n} v_{f\mu_1} \cdots v_{f\mu_n} \sum_{\lambda} \varepsilon'^{(\lambda)*\nu_1 \cdots \nu_n} \varepsilon'^{(\lambda)\mu_1 \cdots \mu_n} . \quad (7)$$

One can show [3] :

$$S_n = \sum_{0 \leq k \leq \frac{n}{2}} C_{n,k} (w_i^2 - 1)^k (w_f^2 - 1)^k (w_i w_f - w_{if})^{n-2k} \quad (8)$$

with $C_{n,k} = (-1)^k \frac{(n!)^2}{(2n)!} \frac{(2n-2k)!}{k!(n-k)!(n-2k)!}$.

From the sum of (4) and (6) one obtains, differentiating relatively to w_{if} [4] ($\ell \geq 0$) :

$$\xi^{(\ell)}(1) = \frac{1}{4} (-1)^\ell \ell! \left\{ \frac{\ell+1}{2\ell+1} 4 \sum_n \left[\tau_{\ell+1/2}^{(\ell)(n)}(1) \right]^2 + \sum_n \left[\tau_{\ell-1/2}^{(\ell-1)(n)}(1) \right]^2 + \sum_n \left[\tau_{\ell-1/2}^{(\ell)(n)}(1) \right]^2 \right\} . \quad (9)$$

This relation shows that $\xi(w)$ is an alternate series in powers of $(w-1)$. Equation (9) reduces to Bjorken SR [1] for $\ell = 1$. Differentiating (6) relatively to w_{if} and making $w_i = w_f = w_{if} = 1$ one obtains :

$$\xi^{(\ell)}(1) = \ell! (-1)^\ell \sum_n \left[\tau_{\ell+1/2}^{(\ell)(n)}(1) \right]^2 \quad (\ell \geq 0) . \quad (10)$$

Combining (9) and (10) one obtains a SR for all ℓ that reduces to Uraltsev SR [2] for $\ell = 1$. From (9) and (10) one obtains :

$$(-1)^\ell \xi^{(\ell)}(1) = \frac{1}{4} \frac{2\ell+1}{\ell} \ell! \left\{ \sum_n \left[\tau_{\ell-1/2}^{(\ell-1)(n)}(1) \right]^2 + \sum_n \left[\tau_{\ell-1/2}^{(\ell)(n)}(1) \right]^2 \right\} . \quad (11)$$

implying

$$\begin{aligned} (-1)^\ell \xi^{(\ell)}(1) &\geq \frac{2\ell+1}{4} \left[(-1)^{\ell-1} \xi^{(\ell-1)}(1) \right] \\ &\geq \frac{(2\ell+1)!!}{2^{2\ell}} \end{aligned} \quad (12)$$

that gives, in particular, for the lower cases,

$$-\xi'(1) = \rho^2 \geq \frac{3}{4} \quad , \quad \xi''(1) \geq \frac{15}{16} \quad (13)$$

Considering systematically the derivatives of the SR (4) and (6) relatively to w_i , w_f , w_{if} with the boundary conditions $w_{if} = w_i = w_f = 1$, one obtains a new SR:

$$\frac{4}{3}\rho^2 + (\rho^2)^2 - \frac{5}{3}\sigma^2 + \sum_{n \neq 0} |\xi^{(n)'}(1)|^2 = 0 \quad (14)$$

that implies :

$$\sigma^2 \geq \frac{1}{5} [4\rho^2 + 3(\rho^2)^2] . \quad (15)$$

There is a simple intuitive argument to understand the term $\frac{3}{5}(\rho^2)^2$ in the best bound (15), namely the non-relativistic quark model, i.e. a non-relativistic light quark q interacting with a heavy quark Q through a potential. The form factor has the simple form :

$$F(\mathbf{k}^2) = \int d\mathbf{r} \varphi_0^+(r) \exp\left(i \frac{m_q}{m_q + m_Q} \mathbf{k} \cdot \mathbf{r}\right) \varphi_0(r) \quad (16)$$

where $\varphi_0(r)$ is the ground state radial wave function. Identifying the non-relativistic IW function $\xi_{NR}(w)$ with the form factor $F(\mathbf{k}^2)$ (16), one can prove that,

$$\sigma_{NR}^2 \geq \frac{3}{5} [\rho_{NR}^2]^2 . \quad (17)$$

Thus, the non-relativistic limit is a good guide-line to study the shape of the IW function $\xi(w)$. We have recently generalized the bound (17) to all the derivatives of $\xi_{NR}(w)$. The method uses the positivity of matrices of moments of the ground state wave function [6]. We have shown that the method can be generalized to the function $\xi(w)$ of QCD.

An interesting phenomenological remark is that the simple parametrization for the IW function [7]

$$\xi(w) = \left(\frac{2}{w+1} \right)^{2\rho^2} \quad (18)$$

satisfies the inequalities (12), (15) if $\rho^2 \geq \frac{3}{4}$.

The result (12), that shows that all derivatives at zero recoil are large, should have important phenomenological implications for the empirical fit needed for the extraction of $|V_{cb}|$ in $B \rightarrow D^* \ell \nu$. The usual fits to extract $|V_{cb}|$ using a linear or linear plus quadratic dependence of $\xi(w)$ are not accurate enough.

A considerable effort has been developed to formulate dispersive constraints on the shape of the form factors in $\bar{B} \rightarrow D^* \ell \nu$ [8]-[9], at finite mass.

Our approach, based on Bjorken-like SR, holds *in the physical region* of the semileptonic decays $\bar{B} \rightarrow D^{(*)} \ell \nu$ and *in the heavy quark limit*. The two approaches are quite different in spirit and in their results.

Let us consider the main results of ref. [9] summarized by the one-parameter formula

$$\xi(w) \cong 1 - 8\rho^2 z + (51\rho^2 - 10)z^2 - (252\rho^2 - 84)z^3 \quad (19)$$

with the variable $z(w)$ defined by

$$z = \frac{\sqrt{w+1} - \sqrt{2}}{\sqrt{w+1} + \sqrt{2}} \quad (20)$$

and the allowed range for ρ^2 being $-0.17 < \rho^2 < 1.51$. This domain is considerably tightened by the lower bound on ρ^2 : $\frac{3}{4} \leq \rho^2 < 1.51$, that shows that our type of bounds are complementary to the bounds obtained from dispersive methods.

2 $1/m_Q$ perturbations of the Current

In this Section, we follow the main lines of our paper [10]. Our starting point is the T -product

$$T_{fi}(q) = i \int d^4x e^{-iq \cdot x} \langle B(p_f) | T[J_f(0) J_i(x)] | B(p_i) \rangle \quad (21)$$

where $J_f(x)$, $J_i(y)$ are the currents (the convenient notation for the subindices i , f will appear clear below) :

$$J_f(x) = \bar{b}(x) \Gamma_f c(x) \quad J_i(y) = \bar{c}(y) \Gamma_i b(y) \quad (22)$$

and p_i is in general different from p_f .

Inserting in this expression hadronic intermediate states, $x^0 < 0$ receives contributions from the direct channel with hadrons with a single heavy quark c , while $x^0 > 0$ receives contributions from hadrons with $b\bar{c}b$ quarks, the Z diagrams :

$$\begin{aligned} T_{fi}(q) = & \sum_{X_c} (2\pi)^3 \delta^3(\mathbf{q} + \mathbf{p}_i - \mathbf{p}_{X_c}) \frac{\langle B_f | J_f(0) | X_c \rangle \langle X_c | J_i(0) | B_i \rangle}{q^0 + E_i - E_{X_c} + i\varepsilon} \\ & - \sum_{X_{\bar{c}bb}} (2\pi)^3 \delta^3(\mathbf{q} - \mathbf{p}_f + \mathbf{p}_{X_{\bar{c}bb}}) \frac{\langle B_f | J_i(0) | X_{\bar{c}bb} \rangle \langle X_{\bar{c}bb} | J_f(0) | B_i \rangle}{q^0 - E_f + E_{X_{\bar{c}bb}} - i\varepsilon} . \quad (23) \end{aligned}$$

We consider the limit $m_c \gg m_b \gg \Lambda_{QCD}$. The difference between the two energy denominators is large $q^0 - E_f + E_{X_{\bar{c}bb}} - (q^0 + E_i - E_{X_c}) \sim 2m_c$ and therefore, we

can in this limit neglect the second term, and we consider the imaginary part of the direct diagram, the first term in (23), the piece proportional to $\delta(q^0 + E_i - E_{X_c})$. Our conditions are, in short, as follows :

$$\Lambda_{QCD} \ll m_b \sim m_c - q^0 \ll q^0 \sim m_c . \quad (24)$$

To summarize, we are considering the heavy quark limit for the c quark, but we allow for a large finite mass for the b quark.

In the conditions (24), or choosing the suitable integration contour, we can write, integrating over q^0

$$T_{fi}^{abs}(\mathbf{q}) \cong \sum_{X_c} (2\pi)^3 \delta^3(\mathbf{q} + \mathbf{p}_i - \mathbf{q}_{X_c}) < B_f | J_f(0) | X_c > < X_c | J_i(0) | B_i > . \quad (25)$$

Finally, integrating over $\mathbf{q}_{X_c}$ and defining $v' = \frac{q+p_i}{m_c}$ one gets

$$T_{fi}^{abs} \cong \sum_{D_n} < B_f(v_f) | J_f(0) | D_n(v') > < D_n(v') | J_i(0) | B_i(v_i) > \quad (26)$$

where we have denoted by $D_n(v')$ the charmed intermediate states.

The T -product matrix element $T_{fi}(q)$ (21) is given, alternatively, in terms of quarks and gluons, by the expression

$$T_{fi}(q) = - \int d^4x e^{-iq \cdot x} < B(p_f) | \bar{b}(0) \Gamma_f S_c(0, x) \Gamma_i b(x) | B(p_i) > \quad (27)$$

where $S_c(0, x)$ is the c quark propagator in the background of the soft gluon field [11].

Since we are considering the absorptive part in the c heavy quark limit of the direct graph in (23), this quantity can be then identified with (27) where $S_c(x, 0)$ is replaced by the following expression [12]

$$S_c(0, x) \rightarrow e^{im_c v' \cdot x} \Phi_{v'}[0, x] D_{v'}(x) \quad (28)$$

where $D_{v'}(x)$ is the *cut* free propagator of a heavy quark

$$D_{v'}(x) = P'_+ \int \frac{d^4k}{(2\pi)^4} \delta(k \cdot v') e^{ik \cdot x} = P'_+ \int_{-\infty}^{\infty} \frac{dt}{2\pi} \delta^4(x - v't) \quad (29)$$

with the positive energy projector defined by $P'_+ = \frac{1}{2}(1 + \not{v}')$.

The eikonal phase $\Phi_{v'}[0, x]$ in (28) corresponds to the propagation of the c quark from the point $x = v't$ to the point 0, that is given by

$$\Phi_{v'}[0, v't] = P \exp \left(-i \int_0^t ds \, v' \cdot A(v's) \right) . \quad (30)$$

This quantity takes care of the dynamics of the soft gluons in HQET along the classical path $x = v't$.

We obtain

$$\begin{aligned} T_{fi}^{abs}(q) &= \int d^4x \, e^{-i(q-m_c v') \cdot x} \int_{-\infty}^{\infty} \frac{dt}{2\pi} \delta^4(x - v't) \\ &< B(p_f) | \bar{b}(0) \Gamma_f P'_+ \Phi_{v'}[0, x] \Gamma_i b(x) | B(p_i) > + O(1/m_c) . \end{aligned} \quad (31)$$

Integrating over x in (31) and making explicit (30),

$$\begin{aligned} T_{fi}^{abs}(q) &= \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{-i(q-m_c v') \cdot v't} \\ &< B(p_f) | \bar{b}(0) \Gamma_f P'_+ P \exp \left(-i \int_0^t ds \, v' \cdot A(v's) \right) \Gamma_i b(v't) | B(p_i) > + O(1/m_c) . \end{aligned} \quad (32)$$

Performing first the integration over q^0 one obtains simply $\delta(v^0 t)$, and making the trivial integration over t one obtains finally the OPE matrix element :

$$T_{fi}^{abs} \cong < B(p_f) | \bar{b}(0) \Gamma_f \frac{1 + \not{v}'}{2v^0} \Gamma_i b(0) | B(p_i) > + O(1/m_c) . \quad (33)$$

Therefore, we end up with the sum rule

$$\begin{aligned} \sum_{D_n} &< B_f(v_f) | J_f(0) | D_n(v') > < D_n(v') | J_i(0) | B_i(v_i) > \\ &= < B(v_f) | \bar{b}(0) \Gamma_f \frac{1 + \not{v}'}{2v^0} \Gamma_i b(0) | B(v_i) > + O(1/m_c) \end{aligned} \quad (34)$$

that is valid for *all powers* of an expansion in $1/m_b$, but only to leading order in $1/m_c$.

On the other hand, making use of the HQET equations of motion, the field $b(x)$ in (34) can be decomposed into upper and lower components as follows [13]

$$b(x) = e^{-im_b v \cdot x} \left(1 + \frac{1}{2m_b + iv \cdot \overrightarrow{D}} i \overrightarrow{\not{D}} \right) h_v(x) \quad (35)$$

where the second term corresponds to the lower components and can be expanded in a series in powers of D_μ/m_b , and v is an arbitrary four-velocity.

Keeping the first order in $1/m_b$, the sum rule reads

$$\begin{aligned}
& \sum_{D_n} \langle B_f(v_f) | J_f(0) | D_n(v') \rangle \langle D_n(v') | J_i(0) | B_i(v_i) \rangle \\
&= \langle B(p_f) | \bar{h}_{v_f}(0) \Gamma_f \frac{1 + \not{v}'}{2v'^0} \Gamma_i h_{v_i}(0) | B(p_i) \rangle \\
&+ \frac{1}{2m_b} \langle B(p_f) | \bar{h}_{v_f}(0) \left[(-i\overleftrightarrow{\not{D}}) \Gamma_f \frac{1 + \not{v}'}{2v'^0} \Gamma_i + \Gamma_f \frac{1 + \not{v}'}{2v'^0} \Gamma_i (i\overrightarrow{\not{D}}) \right] h_{v_i}(0) | B(p_i) \rangle \\
&+ O(1/m_c) + O(1/m_b^2) .
\end{aligned} \tag{36}$$

Therefore, in the OPE side we have, besides the leading dimension 3 operator

$$O^{(3)} = \bar{h}_{v_f} \Gamma_f P'_+ \Gamma_i h_{v_i} \tag{37}$$

the dimension 4 operator

$$O^{(4)} = \bar{h}_{v_f} \left[(-i\overleftrightarrow{\not{D}}) \Gamma_f P'_+ \Gamma_i + \Gamma_f P'_+ \Gamma_i (i\overrightarrow{\not{D}}) \right] h_{v_i} . \tag{38}$$

In the SR we have to compute the l.h.s. including terms of order $1/2m_b$. These terms have been parametrized by Falk and Neubert [14] for the $\frac{1}{2}^-$ doublet and by Leibovich et al. [15] for the transitions between the ground state $\frac{1}{2}^-$ and the $\frac{1}{2}^+, \frac{3}{2}^+$ excited states.

A remark is in order here, that was already made in ref. [17]. Had we taken higher moments of the form $\int dq^0 (q^0)^n T_{fi}^{abs}(q^0)$ ($n > 0$), instead of the lowest one $n = 0$, the integration over q^0 that leads to the simple sum rules (34) or (36) would involve higher dimension operators, giving a whole tower of sum rules [18, 12], even in the *leading* heavy quark limit. Our point of view in this paper is different. We consider the lowest moment $n = 0$, while we expand in powers of $1/m_b$, keeping the first order in this parameter.

Concerning the OPE side in (36), the dimension 4 operator $O^{(4)}$ (38) is nothing else but the $1/m_b$ perturbation of the heavy current $O^{(3)} = \bar{h}_{v_f} \Gamma_f P'_+ \Gamma_i h_{v_i}$ since this operator, containing the Dirac matrix $\Gamma_f P'_+ \Gamma_i$ between heavy quark fields, can be considered as a heavy quark current. Indeed, following Falk and Neubert, the $1/m_b$ perturbation of any heavy quark current $\bar{h}_{v_f} \Gamma h_{v_i}$ is given by

$$\bar{h}_{v_f} \left(-\frac{i\overleftrightarrow{\not{D}}}{2m_b} \right) \Gamma h_{v_i} + \bar{h}_{v_f} \Gamma \left(\frac{i\overrightarrow{\not{D}}}{2m_b} \right) h_{v_i} . \tag{39}$$

However, this perturbation of the current does not exhaust all perturbations in $1/m_b$. We need also to compute the perturbation of the initial and final wave functions $|B_i(v_i)\rangle$, $|B_f(v_f)\rangle$ due to the kinetic and magnetic perturbations of the Lagrangian. This can be done easily following also the prescriptions of Falk and Neubert to compute these corrections in $1/m_b$ for the leading matrix element $\langle B_f(v_f)|\bar{h}_{vf}\Gamma_f P'_+\Gamma_i h_{vi}|B_i(v_i)\rangle$.

The final result is the following. The subleading quantities, functions of w , $\bar{\Lambda}\xi(w)$ and $\xi_3(w)$ [14] can be expressed in terms of leading quantities, namely the $\frac{1}{2}^- \rightarrow \frac{1}{2}^+, \frac{3}{2}^+$ IW functions $\tau_j^{(n)}(w)$ and the corresponding level spacings $\Delta E_j^{(n)}$ ($j = \frac{1}{2}, \frac{3}{2}$) [10]

$$\bar{\Lambda}\xi(w) = 2(w+1) \sum_n \Delta E_{3/2}^{(n)} \tau_{3/2}^{(n)}(1) \tau_{3/2}^{(n)}(w) + 2 \sum_n \Delta E_{1/2}^{(n)} \tau_{1/2}^{(n)}(1) \tau_{1/2}^{(n)}(w) \quad (40)$$

$$\xi_3(w) = (w+1) \sum_n \Delta E_{3/2}^{(n)} \tau_{3/2}^{(n)}(1) \tau_{3/2}^{(n)}(w) - 2 \sum_n \Delta E_{1/2}^{(n)} \tau_{1/2}^{(n)}(1) \tau_{1/2}^{(n)}(w) . \quad (41)$$

These quantities reduce to known SR for $w = 1$, respectively Voloshin SR [16] and a SR for $\xi_3(1)$ [17, 2], and generalizes them for all w .

The comparison of (40), (41) with the results of the BT quark model [7] is very encouraging. Within this scheme $\xi(w)$ is given by (18) with $\rho^2 = 1.02$, while one gets, for the $n = 0$ states

$$\tau_j^{(0)}(w) = \tau_j^{(0)}(1) \left(\frac{2}{w+1} \right)^{2\sigma_j^2} \quad (42)$$

with $\tau_{3/2}^{(0)}(1) = 0.54$, $\sigma_{3/2}^2 = 1.50$, $\tau_{1/2}^{(0)}(1) = 0.22$ and $\sigma_{1/2}^2 = 0.83$. Assuming the reasonable saturation of the SR with the lowest $n = 0$ states [7], one gets, from the first relation (40), a sensibly constant value for $\bar{\Lambda} = 0.513 \pm 0.015$.

3 $1/m_Q$ perturbations of the Lagrangian

We follow closely our recent work [19]. Instead of using the OPE, we will simply use the definition of the subleading elastic $\frac{1}{2}^- \rightarrow \frac{1}{2}^-$ functions $\chi_i(w)$ ($i = 1, 2, 3$) [14]

$$\begin{aligned} \langle D(v')|i \int dx T[J^{cb}(0), \mathcal{L}_v^{(b)}(x)]|B(v)\rangle &= \frac{1}{2m_b} \left\{ -2\chi_1(w) Tr [\bar{D}(v')\Gamma B(v)] \right. \\ &\left. + \frac{1}{2} Tr [A_{\alpha\beta}(v, v')\bar{D}(v')\Gamma P_+ i\sigma^{\alpha\beta} B(v)] \right\} \end{aligned} \quad (43)$$

$$\begin{aligned} < D(v') | i \int dx T[J^{cb}(0), \mathcal{L}_v^{(c)}(x)] | B(v) > = \frac{1}{2m_c} \left\{ -2\chi_1(w) Tr [\bar{D}(v') \bar{\Gamma} B(v)] \right. \\ \left. - \frac{1}{2} Tr [\bar{A}_{\alpha\beta}(v', v) \bar{D}(v') i\sigma^{\alpha\beta} P'_+ \bar{\Gamma} B(v)] \right\} \end{aligned} \quad (44)$$

with

$$\begin{aligned} A_{\alpha\beta}(v, v') &= -2\chi_2(w) (v'_\alpha \gamma_\beta - v'_\beta \gamma_\alpha) + 4\chi_3(w) i\sigma_{\alpha\beta} \\ \bar{A}_{\alpha\beta}(v', v) &= -2\chi_2(w) (v_\alpha \gamma_\beta - v_\beta \gamma_\alpha) - 4\chi_3(w) i\sigma_{\alpha\beta} \end{aligned} \quad (45)$$

where $\bar{A} = \gamma^0 A^+ \gamma^0$ denotes the Dirac conjugate matrix, the current $J^{cb}(0)$ denotes

$$J^{cb} = \bar{h}_v^{(c)} \Gamma h_v^{(b)} \quad (46)$$

where Γ is any Dirac matrix, and $\mathcal{L}_v^{(Q)}(x)$ is given by

$$\mathcal{L}_v^{(Q)} = \frac{1}{2m_Q} [O_{kin,v}^{(Q)} + O_{mag,v}^{(Q)}] \quad (47)$$

with

$$O_{kin,v}^{(Q)} = \bar{h}_v^{(Q)} (iD)^2 h_v^{(Q)} \quad O_{mag,v}^{(Q)} = \frac{g_s}{2} \bar{h}_v^{(Q)} \sigma_{\alpha\beta} G^{\alpha\beta} h_v^{(Q)} . \quad (48)$$

In relations (43)-(45), the $\chi_i(w)$ ($i = 1, 2, 3$) have dimensions of mass, and correspond to the definition given by Luke [20].

We will now insert intermediate states in the T -products (43). We can separately consider $\mathcal{L}_{kin}^{(b)}$ or $\mathcal{L}_{mag}^{(b)}$. The possible Z -diagrams involving heavy quarks contributing to the T -products are suppressed by the heavy quark mass since they are $b\bar{c}c$ intermediate states.

Conveniently choosing the initial and final states, we find the following results :

(1) With $\mathcal{L}_{kin,v}^{(b)}$, pseudoscalar initial state $B(v) = P_+(-\gamma_5)$ and pseudoscalar final state $\bar{D}(v') = \gamma_5 P'_+$, one finds

$$2\chi_1(w) = \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{< B^{(n)}(v) | O_{kin,v}^{(b)}(0) | B(v) >}{\sqrt{4m_{B^{(n)}} m_B}} . \quad (49)$$

In the preceding expressions the energy denominators are $\Delta E_{1/2}^{(n)} = E_{1/2}^{(n)} - E_{1/2}^{(0)}$ ($n \neq 0$).

(2) Consider $\mathcal{L}_{mag,v}^{(b)}$, pseudoscalar initial state $B(v) = P_+(-\gamma_5)$ and pseudoscalar final state $\bar{D}(v') = \gamma_5 P'_+$. Because of parity conservation by the strong interactions, the intermediate states $B^{(n)}$ must have the same parity than the initial state B .

Moreover, $\mathcal{L}_{mag,v}^{(b)}$ being a scalar and producing transitions at zero recoil, the spin of B and $B^{(n)}$ must be the same. Therefore, only pseudoscalar intermediate states $B^{(n)}(0^-)$ can contribute, only states with $j = \frac{1}{2}^-$. One finds, for any current (46)

$$-4(w-1)\chi_2(w) + 12\chi_3(w) = \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{\langle B^{(n)}(v) | O_{mag,v}^{(b)}(0) | B(v) \rangle}{\sqrt{4m_{B^{(n)}}m_B}}. \quad (50)$$

It is remarkable that this linear combination depends only on $\frac{1}{2}^-$ intermediate states.

(3) Consider $\mathcal{L}_{mag,v}^{(b)}$ and a vector initial state $B^*(v, \varepsilon) = P_+ \not{\varepsilon}$ and pseudoscalar final state $\overline{D}(v') = \gamma_5 P_+$. Now we will have vector 1^- intermediate states, either $B^{*(n)}(\frac{1}{2}^-, 1^-)$ or $B^{*(n)}(\frac{3}{2}^-, 1^-)$. For the latter, we have to compute the current matrix element

$$\langle D(v') | J^{cb}(0) | B^{*(n)}(\frac{3}{2}^-, 1^-)(v, \varepsilon) \rangle = \tau_{3/2}^{(2)(n)}(w) Tr [\overline{D}(v') \Gamma F_v^\sigma v'_\sigma] \quad (51)$$

where the $(\frac{3}{2}^-, 1^-)$ operator is given by

$$F_v^\sigma = \sqrt{\frac{3}{2}} P_+ \varepsilon_\nu \left[g^{\sigma\nu} - \frac{1}{3} \gamma^\nu (\gamma^\sigma + v^\sigma) \right] \quad (52)$$

obtained from the $(\frac{3}{2}^+, 1^+)$ operator defined by Leibovich et al. (formula (2.5) of [15]), multiplying by $(-\gamma_5)$ on the right [5]. The Isgur-Wise functions $\tau_{3/2}^{(2)(n)}(w)$ correspond to $\frac{1}{2}^- \rightarrow \frac{3}{2}^-$ transitions, the superindex (ℓ) meaning the orbital angular momentum [4] [5] [10]. As noticed by Leibovich et al. [15], on general grounds the IW functions $\tau_{3/2}^{(2)(n)}(w)$ do not vanish at zero recoil.

One finds, for any current (46), finally

$$\begin{aligned} & -4\chi_2(w)(\varepsilon \cdot v') Tr [\overline{D}(v') \Gamma P_+] + 4\chi_3(w) Tr [\overline{D}(v') \Gamma P_+ \not{\varepsilon}] \\ & = -Tr [\overline{D}(v') \Gamma B^*(v, \varepsilon)] \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{\langle B^{*(n)}(v, \varepsilon) | O_{mag,v}^{(b)}(0) | B^*(v, \varepsilon) \rangle}{\sqrt{4m_{B^{*(n)}}m_{B^*}}} \\ & + \left\{ \sqrt{\frac{3}{2}} (\varepsilon \cdot v') Tr [\overline{D}(v') \Gamma P_+] - \frac{1}{\sqrt{6}} (w-1) Tr [\overline{D}(v') \Gamma P_+ \not{\varepsilon}] \right\} \\ & \sum_n \frac{1}{\Delta E_{3/2}^{(n)}} \tau_{3/2}^{(2)(n)}(w) \frac{\langle B_{3/2}^{*(n)}(v, \varepsilon) | O_{mag,v}^{(b)}(0) | B^*(v, \varepsilon) \rangle}{\sqrt{4m_{B_{3/2}^{*(n)}}m_{B^*}}}. \end{aligned} \quad (53)$$

The energy denominators are $\Delta E_{1/2}^{(n)} = E_{1/2}^{(n)} - E_{1/2}^{(0)}$ ($n \neq 0$) and $\Delta E_{3/2}^{(n)} = E_{3/2}^{(n)} - E_{1/2}^{(0)}$ ($n \geq 0$).

One can obtain other linearly independent relations, taking $\Gamma = \gamma_\mu \gamma_5$.

Since the two four vectors $(v'_\mu - v_\mu)$ and $[(w-1)\varepsilon_\mu + (\varepsilon \cdot v')v_\mu]$ can be chosen to be independent, one obtains independent sum rules for $\chi_2(w)$ and $\chi_3(w)$.

To summarize, making explicit the c flavor, we have obtained the sum rules

$$\chi_1(w) = \frac{1}{2} \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{\langle D^{(n)}(v) | O_{kin,v}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}} m_D}} \quad (54)$$

$$\chi_2(w) = -\frac{3}{4\sqrt{6}} \sum_n \frac{1}{\Delta E_{3/2}^{(n)}} \tau_{3/2}^{(2)(n)}(w) \frac{\langle D_{3/2}^{*(n)}(v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D_{3/2}^{*(n)}} m_{D^*}}} \quad (55)$$

$$\begin{aligned} \chi_3(w) = & -\frac{1}{4} \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{\langle D^{*(n)}(v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}} m_{D^*}}} \\ & - \frac{w-1}{4\sqrt{6}} \sum_n \frac{1}{\Delta E_{3/2}^{(n)}} \tau_{3/2}^{(2)(n)}(w) \frac{\langle D_{3/2}^{*(n)}(v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D_{3/2}^{*(n)}} m_{D^*}}} \end{aligned} \quad (56)$$

There are a number of striking features in relations (54)-(56).

(i) One should notice that *elastic subleading form factors of the Lagrangian type* are given in terms of *leading IW functions*, namely $\xi^{(n)}(w)$ and $\tau_{3/2}^{(2)(n)}(w)$, and *sub-leading form factors at zero recoil*.

(ii) $\chi_1(w)$ is given in terms of matrix elements of $\mathcal{L}_{kin}$, as expected from the definitions (43)-(44) and involve transitions $\frac{1}{2}^- \rightarrow \frac{1}{2}^-$.

(iii) The *elastic subleading magnetic form factors* $\chi_2(w)$ and $\chi_3(w)$ involve $D^*(1^-) \rightarrow D^{*(n)}(1^-)$ transitions $\frac{1}{2}^- \rightarrow \frac{1}{2}^-$ and $\frac{1}{2}^- \rightarrow \frac{3}{2}^-$.

(iv) $\chi_1(w)$ and $\chi_3(w)$ satisfy, as they should, Luke theorem [20],

$$\chi_1(1) = \chi_3(1) = 0 \quad (57)$$

because the $\frac{1}{2}^- \rightarrow \frac{1}{2}^-$ IW functions at zero recoil satisfy

$$\xi^{(n)}(1) = \delta_{n,0} \quad (58)$$

(v) There is a linear combination of $\chi_2(w)$ and $\chi_3(w)$ that gets only contributions from $\frac{1}{2}^- \rightarrow \frac{1}{2}^-$ transitions, namely

$$-4(w-1)\chi_2(w) + 12\chi_3(w) = -3 \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)}(w) \frac{\langle D^{*(n)}(v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}} m_{D^*}}} \quad (59)$$

where the factor -3 is in consistency with (50), shifting from vector to pseudoscalar mesons.

It is well-known that the determination of $|V_{cb}|$ from the $\overline{B} \rightarrow D^* \ell \nu$ differential rate at zero recoil depends on the value of $h_{A_1}(1)$. One interesting point is that precisely the subleading matrix elements of O_{kin} and O_{mag} at zero recoil, that enter in the SR (54)-(56), are related to the quantity $|h_{A_1}(1)|$, as we will see now.

The following SR follows from the OPE [18] [15],

$$\begin{aligned} |h_{A_1}(1)|^2 &+ \sum_n \frac{|\langle D^{*(n)} \left(\frac{1}{2}^-, \frac{3}{2}^- \right) (v, \varepsilon) | \vec{A} | B(v) \rangle|^2}{4m_{D^{*(n)}}m_B} \\ &= \eta_A^2 - \frac{\mu_G^2}{3m_c^2} - \frac{\mu_\pi^2 - \mu_G^2}{4} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) \end{aligned} \quad (60)$$

where $D^{*(n)}$ are 1^- excited states, and

$$\begin{aligned} \mu_\pi^2 &= \frac{1}{2m_B} \langle B(v) | \bar{h}_v^{(b)} (iD)^2 h_v^{(b)} | B(v) \rangle \\ \mu_G^2 &= \frac{1}{2m_B} \langle B(v) | \bar{h}_v^{(b)} \frac{g_s}{2} \sigma_{\alpha\beta} G^{\alpha\beta} h_v^{(b)} | B(v) \rangle \end{aligned} \quad (61)$$

In relation (60) one assumes the states at rest $v = (1, \mathbf{0})$ and the axial current is space-like, orthogonal to v .

In the l.h.s. of relation (60),

$$h_{A_1}(1) = \eta_{A_1} + \delta_{1/m^2}^{(A_1)} \quad (62)$$

($\eta_{A_1} = 1$ + radiative corrections) because there are no first order $1/m_Q$ corrections due to Luke theorem. The sum over the squared matrix elements of $B \rightarrow D^{*(n)}(1^-)$ transitions contains two types of possible contributions, corresponding to $D^{*(n)} \left(\frac{1}{2}^-, 1^- \right)$ ($n \neq 0$), and $D^{*(n)} \left(\frac{3}{2}^-, 1^- \right)$ ($n \geq 0$). The r.h.s. of (60) exhibits the OPE at the desired order. From the decomposition between radiative corrections and $1/m_Q^2$ corrections (62) one gets, from (60), neglecting higher order terms,

$$\begin{aligned} -\delta_{1/m^2}^{(A_1)} &= \frac{\mu_G^2}{6m_c^2} + \frac{\mu_\pi^2 - \mu_G^2}{8} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) \\ &+ \frac{1}{2} \sum_n \frac{|\langle D^{*(n)} \left(\frac{1}{2}^-, \frac{3}{2}^- \right) (v, \varepsilon) | \vec{A} | B(v) \rangle|^2}{4m_{D^{*(n)}}m_B}. \end{aligned} \quad (63)$$

The correction $\delta_{1/m^2}^{(A_1)}$ is therefore negative, both terms being of the same sign.

The matrix elements $\langle D^{*(n)} \left(\frac{1}{2}^-, \frac{3}{2}^- \right) (v, \varepsilon) | \vec{A} | B \rangle$ have been expressed in terms of the matrix elements $\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{kin,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle$ and $\langle D^{*(n)} \left(\frac{1}{2}^-, \frac{3}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle$ by Leibovich et al. (formulas (4.1) and (4.3)) [15]. Hence, $-\delta_{1/m^2}^{(A_1)}$ (63) can be written as

$$\begin{aligned}
-\delta_{1/m^2}^{(A_1)} = & \frac{\mu_G^2}{6m_c^2} + \frac{1}{8} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) (\mu_\pi^2 - \mu_G^2) \\
& + \frac{1}{2} \sum_n \left[\left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) \frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right. \\
& + \left. \left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) \frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{kin,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right]^2 \\
& + \frac{1}{2} \sum_n \left[\frac{1}{2m_c} \frac{1}{\Delta E_{3/2}^{(n)}} \frac{\langle D^{*(n)} \left(\frac{3}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right]^2. \quad (64)
\end{aligned}$$

The important point to emphasize here is that the matrix elements $\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{kin,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle$ and $\langle D^{*(n)} \left(\frac{1}{2}^-, \frac{3}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle$ are precisely the same ones that enter in the SR (54)-(56). This allows to obtain an interesting lower bound on $-\delta_{1/m^2}^{(A_1)}$.

Taking now the relevant linear combinations of the matrix elements suggested by the r.h.s. of (64), using (54), (55) and (59), and Schwarz inequality

$$\left| \sum_n A_n B_n \right| \leq \sqrt{\left(\sum_n |A_n|^2 \right) \left(\sum_n |B_n|^2 \right)} \quad (65)$$

one finds

$$\begin{aligned}
& \sum_{n \neq 0} \left[\xi^{(n)}(w) \right]^2 \sum_{n \neq 0} \left\{ \frac{1}{\Delta E_{1/2}^{(n)}} \left[\left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) \frac{\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{kin,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right. \right. \\
& + \left. \left. \left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) \frac{\langle D^{*(n)} \left(\frac{1}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right] \right\}^2 \\
& \geq 4 \left\{ \left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) \chi_1(w) - \frac{1}{3} \left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) [-2(w-1)\chi_2(w) + 6\chi_3(w)] \right\}^2 \quad (66)
\end{aligned}$$

$$\begin{aligned}
& \sum_n \left[\tau_{3/2}^{(2)(n)}(w) \right]^2 \sum_n \left\{ \frac{1}{\Delta E_{3/2}^{(n)}} \left[\frac{1}{2m_c} \frac{\langle D^{*(n)} \left(\frac{3}{2}^- \right) (v, \varepsilon) | O_{mag,v}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{*(n)}}m_{D^*}}} \right] \right\}^2 \\
& \geq \frac{32}{3} \left[\frac{1}{2m_c} \chi_2(w) \right]^2. \quad (67)
\end{aligned}$$

These two last equations imply, from (64), the inequality

$$\begin{aligned}
-\delta_{1/m^2}^{(A_1)} &\geq \frac{\mu_G^2}{6m_c^2} + \frac{\mu_\pi^2 - \mu_G^2}{8} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) \\
&+ 2 \frac{\left\{ \left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) \chi_1(w) - \frac{1}{3} \left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) [-2(w-1)\chi_2(w) + 6\chi_3(w)] \right\}^2}{\sum_{n \neq 0} [\xi^{(n)}(w)]^2} \\
&+ \frac{16}{3} \frac{\left[\frac{1}{2m_c} \chi_2(w) \right]^2}{\sum_n [\tau_{3/2}^{(2)(n)}(w)]^2} . \tag{68}
\end{aligned}$$

This inequality on $-\delta_{1/m^2}^{(A_1)}$ involves on the r.h.s. *elastic subleading* functions $\chi_i(w)$ ($i = 1, 2, 3$) in the numerator and sums over *inelastic leading IW functions* $\sum_{n \neq 0} [\xi^{(n)}(w)]^2$ and $\sum_n [\tau_{3/2}^{(2)(n)}(w)]^2$ in the denominator. We must emphasize that this inequality is valid for all values of w and constitutes a rigorous constraint between these functions and the correction $-\delta_{1/m^2}^{(A_1)}$. Let us point out that, near $w = 1$, since $\xi^{(n)}(w) \sim (w-1)$ ($n \neq 0$) and, due to Luke theorem $\chi_1(w)$, $\chi_3(w) \sim (w-1)$, the second term on the r.h.s. of (68) is a constant in the limit $w \rightarrow 1$.

On the other hand, since $\chi_2(w)$ is not protected by Luke theorem, $\chi_2(1) \neq 0$ and in general, as pointed out by Leibovich et al. $\tau_{3/2}^{(2)}(1) \neq 0$, the last term in the r.h.s. of (68) is also a constant for $w = 1$.

The inequality (68) is valid for all values of w , and in particular it holds in the $w \rightarrow 1$ limit. Let us consider this limit, that gives

$$\begin{aligned}
-\delta_{1/m^2}^{(A_1)} &\geq \frac{\mu_G^2}{6m_c^2} + \frac{\mu_\pi^2 - \mu_G^2}{8} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) \\
&+ 2 \frac{\left\{ \left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) \chi'_1(1) - \frac{1}{3} \left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) [-2\chi_2(1) + 6\chi'_3(1)] \right\}^2}{\sum_{n \neq 0} [\xi^{(n)'}(1)]^2} \\
&+ \frac{16}{3} \frac{\left[\frac{1}{2m_c} \chi_2(1) \right]^2}{\sum_n [\tau_{3/2}^{(2)}(1)]^2} . \tag{69}
\end{aligned}$$

On the other hand, using the OPE in the heavy quark limit, we have demonstrated above the following sum rules [4]

$$\sum_n [\tau_{3/2}^{(2)}(1)]^2 = \frac{4}{5} \sigma^2 - \rho^2 \tag{70}$$

$$\sum_{n \neq 0} [\xi^{(n)'}(1)]^2 = \frac{5}{3} \sigma^2 - \frac{4}{3} \rho^2 - (\rho^2)^2 \tag{71}$$

where ρ^2 and σ^2 are the slope and the curvature of the elastic Isgur-Wise function $\xi(w)$.

The positivity of the l.h.s. of (70), (71) yield respectively the lower bounds on the curvature obtained (13) and (15). Relations (69)-(71) give finally the bound

$$\begin{aligned}
-\delta_{1/m^2}^{(A_1)} &\geq \frac{\mu_G^2}{6m_c^2} + \frac{\mu_\pi^2 - \mu_G^2}{8} \left(\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_cm_b} \right) \\
&+ \frac{2}{3[5\sigma^2 - 4\rho^2 - 3(\rho^2)^2]} \left\{ \left(\frac{1}{2m_c} - \frac{1}{2m_b} \right) 3\chi'_1(1) - \left(\frac{1}{2m_c} + \frac{3}{2m_b} \right) [-2\chi_2(1) + 6\chi'_3(1)] \right\}^2 \\
&+ \frac{80}{3(4\sigma^2 - 5\rho^2)} \left[\frac{1}{2m_c} \chi_2(1) \right]^2. \tag{72}
\end{aligned}$$

A number of remarks are worth to be made here :

(i) The bounds contain an OPE piece, dependent on μ_π^2 and μ_G^2 , and a piece that bounds the inelastic contributions, given in terms of the $1/m_Q$ elastic quantities $\chi'_1(1)$, $\chi_2(1)$, $\chi'_3(1)$ and of the slope ρ^2 and curvature σ^2 of the elastic IW function $\xi(w)$.

(ii) Taking roughly constant values for $\chi'_1(1)$, $\chi_2(1)$, $\chi'_3(1)$, independent of ρ^2 and σ^2 , as suggested by the QCD Sum Rules calculations (QCDSR) [21] [22] [23], the bounds for the inelastic contributions diverge in the limit $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$, according to (13). This feature does not seem to us physical.

(iii) *Therefore, one should expect that $\chi'_1(1)$, $\chi_2(1)$ and $\chi'_3(1)$ vanish also in this limit.* We give a demonstration of this interesting feature below.

(iv) Thus, the limit $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$ seems related to the behaviour of $\chi_i(w)$ ($i = 1, 2, 3$) near zero recoil.

(v) The feature (iii) does not appear explicitly in the QCDSR approach, where one gets roughly $\rho_{ren}^2 \cong 0.7$, and where there is no dependence on ρ^2 of the functions $\chi_i(w)$ ($i = 1, 2, 3$).

Now we demonstrate that indeed $\chi'_1(1)$, $\chi_2(1)$ and $\chi'_3(1)$ vanish in the limit $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$. At zero recoil $w \rightarrow 1$ we have

$$\chi'_1(1) = \frac{1}{2} \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)'}(1) \frac{\langle D^{(n)}(v) | O_{kin}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \tag{73}$$

$$\chi_2(1) = \frac{1}{4\sqrt{6}} \sum_n \frac{1}{\Delta E_{3/2}^{(n)}} \tau_{3/2}^{(2)(n)}(1) \frac{\langle D_{3/2}^{*(n)}(v, \varepsilon) | O_{mag}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D_{3/2}^{*(n)}}m_{D^*}}} \tag{74}$$

$$-4\chi_2(1) + 12\chi_3'(1) = \sum_{n \neq 0} \frac{1}{\Delta E_{1/2}^{(n)}} \xi^{(n)'}(1) \frac{\langle D^{(n)}(v) | O_{mag}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \quad (75)$$

Using again Schwarz inequality, we obtain

$$[\chi_1'(1)]^2 \leq \frac{1}{4} \sum_{n \neq 0} [\xi^{(n)'}(1)]^2 \sum_{n \neq 0} \left[\frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{(n)}(v) | O_{kin}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \right]^2 \quad (76)$$

$$[\chi_2(1)]^2 \leq \frac{1}{96} \sum_n [\tau_{3/2}^{(2)(n)}(1)]^2 \sum_n \left[\frac{1}{\Delta E_{3/2}^{(n)}} \frac{\langle D_{3/2}^{*(n)}(v, \varepsilon) | O_{mag}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{(n)}}m_{D^*}}} \right]^2 \quad (77)$$

$$[-4\chi_2(1) + 12\chi_3'(1)]^2 \leq \sum_{n \neq 0} [\xi^{(n)'}(1)]^2 \sum_{n \neq 0} \left[\frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{(n)}(v) | O_{mag}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \right]^2 \quad (78)$$

and from relations (70) and (71),

$$[\chi_1'(1)]^2 \leq \frac{1}{12} [5\sigma^2 - 4\rho^2 - 3(\rho^2)^2] \sum_{n \neq 0} \left[\frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{(n)}(v) | O_{kin}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \right]^2 \quad (79)$$

$$[\chi_2(1)]^2 \leq \frac{1}{480} (4\sigma^2 - 5\rho^2) \sum_n \left[\frac{1}{\Delta E_{3/2}^{(n)}} \frac{\langle D_{3/2}^{*(n)}(v, \varepsilon) | O_{mag}^{(c)}(0) | D^*(v, \varepsilon) \rangle}{\sqrt{4m_{D^{(n)}}m_{D^*}}} \right]^2 \quad (80)$$

$$\begin{aligned} & [-4\chi_2(1) + 12\chi_3'(1)]^2 \\ & \leq \frac{1}{3} [5\sigma^2 - 4\rho^2 - 3(\rho^2)^2] \sum_{n \neq 0} \left[\frac{1}{\Delta E_{1/2}^{(n)}} \frac{\langle D^{(n)}(v) | O_{mag}^{(c)}(0) | D(v) \rangle}{\sqrt{4m_{D^{(n)}}m_D}} \right]^2 \end{aligned} \quad (81)$$

Therefore, in the limit $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$, one obtains

$$\chi_1'(1) = \chi_2(1) = \chi_3'(1) = 0 \quad (82)$$

This is a very strong correlation relating the behaviour of the elastic IW function $\xi(w)$ to the elastic subleading IW functions $\chi_i(w)$ ($i = 1, 2, 3$) near zero recoil.

To conclude, we have obtained bounds that relate the $1/m_Q^2$ correction of the form factor $h_{A_1}(w)$ to the $1/m_Q$ subleading form factors of the Lagrangian type $\chi_i(w)$ ($i = 1, 2, 3$) and to the shape of the elastic Isgur-Wise $\xi(w)$. This bound should in principle be taken into account in the analysis of the exclusive determination of $|V_{cb}|$ in the channels $\overline{B} \rightarrow D(D^*)\ell\nu$. On the other hand, we have demonstrated an

important constraint on the behavior of the subleading form factors $\chi_i(w)$: in the limit $\rho^2 \rightarrow \frac{3}{4}$, $\sigma^2 \rightarrow \frac{15}{16}$, $\chi_1'(1)$, $\chi_2(1)$ and $\chi_3'(1)$ must vanish.

It would be very interesting to have a theoretical calculation of the functions $\chi_i(w)$ ($i = 1, 2, 3$) satisfying this constraint. Otherwise it seems questionable to try an exclusive determination of $|V_{cb}|$ by fitting the slope ρ^2 and considering uncorrelated subleading corrections.

In conclusion, using sum rules in HQET, as formulated in ref. [3, 4, 10, 19], we have found lower bounds for the moduli of the derivatives of $\xi(w)$ and non-trivial results on $1/m_Q$ form factors of the Current and Lagrangian types. We have also obtained a lower bound on $-\delta_{1/m^2}^{(A_1)}$. The determination of the CKM matrix element $|V_{cb}|$ in $B \rightarrow D^{(*)}\ell\nu$ should satisfy these constraints.

Acknowledgement

We are indebted to the EC contract HPRN-CT-2002-00311 (EURIDICE) for its support.

References

- [1] J. D. Bjorken, invited talk at Les Rencontres de la Vallée d'Aoste, La Thuile, SLAC-PUB-5278, 1990 ; N. Isgur and M. B. Wise, *Phys. Rev. D* **43**, 819 (1991).
- [2] N. Uraltsev, *Phys. Lett. B* **501**, 86 (2001).
- [3] A. Le Yaouanc, L. Oliver and J.-C. Raynal, *Phys. Rev. D* **67**, 114009 (2003).
- [4] A. Le Yaouanc, L. Oliver and J.-C. Raynal, *Phys. Lett. B* **557**, 207 (2003).
- [5] A. Falk, *Nucl. Phys. B* **378**, 79 (1992).
- [6] F. Jugeau, A. Le Yaouanc, L. Oliver and J.-C. Raynal, hep-ph/0405234, *Phys. Rev. D* **70**, 114020 (2004).
- [7] V. Morénas, A. Le Yaouanc, L. Oliver, O. Pène and J.-C. Raynal, *Phys. Rev. D* **56**, 5668 (1997).
- [8] C. G. Boyd, B. Grinstein and R. F. Lebed, *Phys. Rev. D* **56**, 6895 (1997).

- [9] I. Caprini, L. Lellouch and M. Neubert, *Nucl. Phys. B* **530**, 153 (1998).
- [10] J. Jugeau, A. Le Yaouanc, L. Oliver and J.-C. Raynal, hep-ph/0407176, *Phys. Rev. D* **71**, 054031 (2005).
- [11] B. Blok, L. Koryakh, M. Shifman and A. Vainshtein, *Phys. Rev. D* **49**, 3356 (1994).
- [12] A. Grozin and G. Korchemsky, *Phys. Rev. D* **53**, 1378 (1996).
- [13] See, for example, M. Neubert, Lectures at the Trieste Summer School in Particle Physics (Part II) (1999), hep-ph/0001334.
- [14] A. Falk and M. Neubert, *Phys. Rev. D* **47**, 2965 (1993).
- [15] A. Leibovich et al., *Phys. Rev. D* **57**, 308 (1998).
- [16] M. Voloshin, *Phys. Rev. D* **46**, 3062 (1992).
- [17] A. Le Yaouanc et al., *Phys. Lett. B* **480**, 119 (2000).
- [18] I. Bigi et al., *Phys. Rev. D* **52**, 196 (1995).
- [19] F. Jugeau, A. Le Yaouanc, L. Oliver and J.-C. Raynal, LPT Orsay 05-43, hep-ph/0510178.
- [20] M. E. Luke, *Phys. Lett. B* **252**, 447 (1990).
- [21] M. Neubert, *Phys. Rev. B* **46**, 3914 (1992).
- [22] M. Neubert, Z. Ligeti and Y. Nir, *Phys. Lett. B* **301**, 101 (1993) ; *Phys. Rev. D* **47**, 5060 (1993).
- [23] M. Neubert, *Phys. Reports* **245**, 259 (1994).